

High-probability quantum state transfer among nodes of an open XXZ spin chain

E.B.Fel'dman, A.I. Zenchuk *

*Institute of Problems of Chemical Physics, Russian Academy of Sciences,
Chernogolovka, Moscow reg., 142432, Russia*

Abstract

This paper concerns the problem of the high probability state transfer among 2^s symmetrically placed nodes of the N -nodes spins $1/2$ chain with the XXZ Hamiltonian. We consider examples with $(N, s) = (4, 4)$, $(N, s) = (6, 4)$ and $(N, s) = (8, 8)$.

Key words: spin dynamics, quantum state transfer, spin chain, ideal state transfer, high-probability state transfer

PACS: 05.30.-d, 76.20.+q

1 Introduction

This paper concerns the problem of the high probability state transfer (HPST) [1] between different nodes of the spin $1/2$ chain, which becomes a popular problem due to the development of the quantum communication systems and quantum computing. By "state transfer" we mean the following phenomenon [2,3]. Consider the chain of spins $1/2$ with dipole-dipole interactions in the strong external magnetic field. Let all spins be directed along the external magnetic field except the i th one whose initial state is arbitrary, $\psi_i = \cos \theta |0\rangle + e^{i\phi} \sin \theta |1\rangle$, where $|0\rangle$ and $|1\rangle$ mean the spin directed along and opposite the external magnetic field respectively. Let the energy of the ground state (all spins are aligned along the magnetic field) be zero. If the state of j th node becomes $\psi_j = \cos \theta |0\rangle + e^{i\phi} f_{ij} \sin \theta |1\rangle$ with $|f_{ij}| = 1$ at time moment t_0 then we say that the state has been transferred from the i th to the j th node with the phase shift $\phi_{ij} = \arg f_{ij}$. Since $|f_{ij}| = 1$, all other spins are directed along the

* Corresponding author.

Email addresses: efeldman@icp.ac.ru (E.B.Fel'dman), zenchuk@itp.ac.ru (A.I. Zenchuk).

field at $t = t_0$, i.e. their states are $|0\rangle$. Function f_{ij} is the transition amplitude of an excited state $|1\rangle$ from the i th to the j th node. Note that if all nodes of the chain have equal Larmor frequencies and we are interested in the state propagation between two nodes, say between s th to r th nodes, then the shift ϕ_{sr} may be simply removed by the proper choice of the constant magnetic field value [2], so that $f_{ij} = 1$, i.e. the state is perfectly transferred.

There exists a wide literature studying the state transfer along the spin chains in the strong external magnetic field. For instance, propagation of the spin waves in homogeneous chains was considered in [4]. It was demonstrated that the state may be perfectly transferred between two end nodes [3,5] as well as between two symmetrical inner nodes [6] in the inhomogeneous chain. State propagation along the alternating chains was studied in [7,8]. It was shown in [9,10] that the chains with weak end bonds provide the state transfer from one to another side of the chain. End-to-end entanglement in both alternating chains and chains with weak end bonds has been studied in [9]. Some aspects of the entanglement between remote nodes of the chain have been studied in [11].

However, all these references consider the state transfer between two end nodes (or between two symmetrical nodes) which is required for the construction of the communication channels where the state must be transferred from one object to another. Meanwhile, the quantum computation requires such systems which have HPSTs among many different nodes and, as a consequence, may distribute information among these nodes. Such systems may be candidates for the quantum register. Emphasise that, as it was indicated above, the excited state may be transferred from the initial i th node to some j th node with proper phase shift ϕ_{ij} . However, we will show that all these shifts may be removed in a simple way introducing the time dependent magnetic field, see Sec.2 (remember that the single phase shift can be removed by a constant magnetic field, like it was done in the case of the state propagation between two nodes [2]). Thus, in general, the phase shifts ϕ_{ij} do not create serious obstacles for the quantum communications. The only problem is the organization of the state transfers with big values of $|f_{ij}|$.

A simple variant of such systems is suggested in our paper. Namely, we construct the chain of N nodes which has set $\mathcal{L}$ of $\mathcal{N} \leq N$ nodes p_i , $i = 1, \dots, \mathcal{N}$,

$$\mathcal{L} = \{p_i, \ i = 1, \dots, \mathcal{N} \leq N\}, \quad p_1 \equiv 1, \quad (1)$$

with the HPSTs between any two of them, i.e. if the unknown state $\phi_n = \cos \theta |0\rangle + e^{i\phi} \sin \theta |1\rangle$ is generated in any particular node n from the set $\mathcal{L}$ (while the initial states of all other nodes are $|0\rangle$) then this state may be detected with high probability in any other node m from the set $\mathcal{L}$ after appropriate time intervals.

Hereafter we will use the notation $L_{1,\dots,K}(N_1, M_1, N_2, M_2, \dots, N_K - 1, M_K - 1, N_K)$ for the chain shown in Fig.1. Here N_i and M_i are the numbers of nodes




Fig. 1. The non-symmetric chain $L_{1,\dots,K}(N_1, M_1, N_2, M_2, \dots, N_K - 1, M_K - 1, N_K)$. The total number of the nodes is $N = \sum_{i=1}^{K-1} (N_i + M_i) + N_K$. Here $\mathcal{L} = \{p_i, i = 1, \dots, K\}$. The chains L_i allow the HPSTs among their end nodes p_{2i-1} and p_{2i} , $i = 1, \dots, K$. These chains are connected through the spin 1/2 chains C_i , $i = 1, \dots, K - 1$. Parameters $\mathcal{D}_{i(2i)}$ ($\mathcal{D}_{i(2i+1)}$) are coupling constants between p_{2i} (p_{2i+1}) and first (last) node of the chain C_i , $i = 1, \dots, K - 1$. For unambiguity, the coupling constants between n th and m th nodes of the whole chain will be called D_{nm} . Namely parameters D_{nm} appear in the Hamiltonian (12). $D_n \equiv D_{n(n+1)}$ are coupling constants between the nearest neighbours.

in L_i and C_i respectively. Let us clarify the structure of this scheme. Each of the chains L_i , $i = 1, \dots, K$, allows the HPST between its end nodes p_{2i-1} and p_{2i} . Chains l_i collect all inner nodes of L_i and consequently have $N_i - 2$ nodes. Two chains L_i and L_{i+1} are connected by the "weak bond" through the chain C_i , $i = 1, \dots, K - 1$. By "weak bond" between L_i and L_{i+1} we mean the following necessary inequality among the coupling constants:

$$\max(\mathcal{D}_{i(2i)}, \mathcal{D}_{i(2i+1)}, \text{coupling constants in } C_i, i = 1, \dots, K - 1) < \min(\text{coupling constants in } L_i, i = 1, \dots, K). \quad (2)$$

Emphasize that, also the HPST is organized between end nodes of each particular chain L_i (taken out of the general chain), the whole chain $L_{1,\dots,K}$ does not provide the HPSTs between all p_i ($i = 1, \dots, 2K$) in general case. However, we are interested in the particular form of the chain $L_{1,\dots,K}$ which does provide the HPST among all nodes p_i . First of all, such chain must be symmetrical and may be written as ($K = \mathcal{N}/2$)

$$L_{1,\dots,\mathcal{N}/4-1,\mathcal{N}/4,\mathcal{N}/4,\mathcal{N}/4-1,\dots,1}(N_1, M_1, \dots, N_{\mathcal{N}/4}, M_{\mathcal{N}/4}, N_{\mathcal{N}/4}, \dots, M_1, N_1),$$

where $\mathcal{N} = 2^s$, $s = 1, 2, \dots$, see Fig.2.




Fig. 2. General scheme of the symmetric spin 1/2 chain with the HPSTs among the nodes p_i , $i = 1, \dots, \mathcal{N}$, $\mathcal{N} = 2^s$, $s = 1, 2, \dots$. Here the total number of the nodes $N = 2 \sum_{i=1}^{\mathcal{N}/4-1} (N_i + M_i) + 2N_{\mathcal{N}/4} + M_{\mathcal{N}/4}$, $\mathcal{L} = \{p_i, i = 1, \dots, \mathcal{N}\}$.

Let the chain have N nodes. We use notations $\bar{P}_{p_n p_m}^{(N)}$, $\bar{t}_{p_n p_m}^{(N)}$ and $\bar{\phi}_{p_n p_m}^{(N)}$ for the probability of the exited state transfer between p_n th and p_m th nodes, for the time interval required for this transfer and for the phase shift of the transfered exited state, $n, m = 1, \dots, \mathcal{N} \leq N$:

$$\bar{P}_{p_n p_m}^{(N)} \equiv P_{p_n p_m}^{(N)}(\bar{t}_{p_n p_m}^{(N)}) = |f_{p_i p_j}(\bar{t}_{p_n p_m}^{(N)})|^2, \quad \bar{\phi}_{p_i p_j}^{(N)} = \arg f_{p_i p_j}(\bar{t}_{p_n p_m}^{(N)}). \quad (3)$$

Due to the symmetry of the chain, we have the following identities:

$$\begin{aligned} \bar{P}_{p_n p_m}^{(N)} &= \bar{P}_{p_m p_n}^{(N)}, \quad \bar{t}_{p_n p_m}^{(N)} = \bar{t}_{p_m p_n}^{(N)}, \quad \bar{\phi}_{p_n p_m}^{(N)} = \bar{\phi}_{p_m p_n}^{(N)} \\ \bar{P}_{p_n p_m}^{(N)} &= \bar{P}_{(N-p_n+1)(N-p_m+1)}^{(N)}, \quad \bar{t}_{p_n p_m}^{(N)} = \bar{t}_{(N-p_n+1)(N-p_m+1)}^{(N)}, \\ \bar{\phi}_{p_n p_m}^{(N)} &= \bar{\phi}_{(N-p_n+1)(N-p_m+1)}^{(N)}. \end{aligned} \quad (4)$$

Because of the wide spread of the coupling constants, the time interval $\bar{t}_{p_n p_m}^{(N)}$ needed for the state transfer between p_n and p_m significantly depends on the values n and m . In general,

$$\bar{t}_{p_n p_m}^{(N)} \big|_{n \leq \mathcal{N}/2, m > \mathcal{N}/2} \ll \bar{t}_{p_n p_m}^{(N)} \big|_{n, m \leq \mathcal{N}/2}, \quad (5)$$

i.e. the state may be transfered between two nodes much faster if both nodes are placed in the same half of the chain. Thus, an important characteristic of such chain is the interval

$$T_{\mathcal{N}}^{(N)} = \max \left(\bar{t}_{p_n p_m}^{(N)}, \quad n, m = 1, \dots, \mathcal{N} \right). \quad (6)$$

Hereafter the state transfer between the nodes p_n and p_m will be referred to as HPST if

$$\bar{P}_{p_n p_m}^{(N)} \geq P_0. \quad (7)$$

The value P_0 is conventional. We take $P_0 = 0.9$ in Examples of Sec.3 and in Example 1 of Sec.5 and $P_0 = 0.8$ in Examples of Sec.4 and in Example 2 of Sec.5.

The set of all possible HPSTs between any two nodes from the list $\mathcal{L}$ will be referred to as HPST($N; \mathcal{L}$), where N is the total number of nodes in the chain. We call the parameters of HPST($N; \mathcal{L}$) in such chain the set of parameters

$$\bar{P}_{p_n p_m}^{(N)}, \quad \bar{t}_{p_n p_m}^{(N)}, \quad \bar{\phi}_{p_n p_m}^{(N)}. \quad (8)$$

Since the organization of the HPSTs among different nodes of the set of $\mathcal{N} > 2$ nodes is an essential property of the quantum register, the chains constructed in this paper may be candidates for this role.

While the nodes p_i may serve as the q-bits of the quantum register, the chains l_i (and C_i) serve to decrease the time intervals $\bar{t}_{p_{2i-1}p_{2i}}^{(N)}$ (and $\bar{t}_{p_{2i}p_{2i+1}}^{(N)}$) required for the state transfer between the nodes p_{2i-1} and p_{2i} (and between the nodes p_{2i} and p_{2i+1}) separated by the long distance as it happens in the communication channels. Namely, if L_i consists of two nodes p_{2i-1} and p_{2i} then the time interval $\bar{t}_{p_{2i-1}p_{2i}}^{(N)}$ may be reduced putting additional chain l_i with properly adjusted coupling constants between these two nodes. Similarly, the time interval $\bar{t}_{p_{2i}p_{2i+1}}^{(N)}$ required to transfer the excited state between the last node of L_i (i.e. node p_{2i}) and the first node of L_{i+1} (i.e. node p_{2i+1}) may be reduced putting chain C_i with properly adjusted coupling constants between L_i and L_{i+1} , see, for instance, chain $C_1(2)$ in Fig.8. Thus the above spin chain $L_{1,\dots,\mathcal{N}/4-1,\mathcal{N}/4,\mathcal{N}/4-1,\dots,1}(N_1, M_1, \dots, N_{\mathcal{N}/4}, M_{\mathcal{N}/4}, N_{\mathcal{N}/4}, \dots, M_1, N_1)$ combines properties of both quantum register and communication channel.

Finally we note that, constructing the spin chain, we want

$$(1) \text{ to satisfy the condition (7) for } \bar{P}_{p_n p_m}^{(N)}, n, m = 1, \dots, \mathcal{N}, \quad (9)$$

$$(2) \text{ to minimize the parameter } T_{\mathcal{N}}^{(N)}, \quad (10)$$

$$(3) \text{ to remove phase shifts } \bar{\phi}_{p_n p_m}^{(N)}, n, m = 1, \dots, \mathcal{N}. \quad (11)$$

This paper is organized as follows. In Sec.2 we show that $2^N \times 2^N$ matrix representation of the XXZ Hamiltonian may be reduced to the $N \times N$ matrix representation for a certain type of problems. Sec.3 describes the general structures of the spin chains having the set $\mathcal{L}$ of four nodes; simple examples are represented. Similar study of the eight node chain is given in Sec.4. Deformations of the above chains decreasing parameter $T_{\mathcal{N}}^{(N)}$ are discussed in Sec.5. Conclusions are given in Sec.6.

2 State transfer in spin 1/2 chain with XXZ Hamiltonian

We study the HPSTs [1] among nodes of the spin 1/2 chain in strong external magnetic field $B(t)$ with the XXZ Hamiltonian

$$\mathcal{H} = \mathcal{H}_{dz} + \omega(t)I_z, \quad I_z = \sum_{i=1}^N I_{i,z}, \quad \omega(t) = \gamma B(t) \quad (12)$$

$$\mathcal{H}_{dz} = \sum_{\substack{i,j=1 \\ j>i}}^N D_{ij}(I_{i,x}I_{j,x} + I_{i,y}I_{j,y} - 2I_{i,z}I_{j,z}), \quad D_{ij} = \frac{\gamma^2 \hbar}{r_{ij}^3},$$

where γ is the gyromagnetic ratio, r_{ij} is the distance between i th and j th spins, $I_{i,\alpha}$ is the projection operator of the i th spin on the α axis, $\alpha = x, y, z$, I_z is the z -projection operator of the total spin, D_{ij} are the spin-spin coupling constants. This Hamiltonian describes the secular part of the dipole-dipole interaction in the strong external magnetic field [12]. For our convenience, we call $D_i \equiv D_{i(i+1)}$, $i = 1, \dots, N-1$. It is obvious that all D_i may be arbitrary by definition.

Although the approximation of the Hamiltonian (12) by the nearest neighbour interaction is very popular in study of the quantum state transfer, one can show that it is not applicable to the inhomogeneous chains which have a wide spread of coupling constants. In fact, this approximation is applicable if $D_{ij} \ll \min(D_k, k = 1, \dots, N-1)$, $\forall j > i+1$. Only in this case we may disregard terms with coupling constants D_{ij} ($j > i+1$) in the Hamiltonian. However, this is not possible in general. Consider, for instance, the chain in Fig.3. In this chain, the approximation by the nearest neighbour interaction




Fig. 3. Example of the chain which does not allow the approximation by the nearest neighbour interaction

takes into account the term with coupling constant $D_{i+2} = \frac{\alpha}{64a^3}$ and disregards the term with coupling constant $D_{i(i+2)} = \frac{\alpha}{27a^3} > D_{i+2}$ in the Hamiltonian, i.e. we neglect the term which is bigger then the term which is taken into account. Comparison of the coupling constants in the chains considered in Secs.3-5 confirms that the approximation of the Hamiltonian by the nearest neighbour interaction is not applicable to these chains. Thus, hereafter we consider the total Hamiltonian (12).

It is convenient to take the eigenvectors of I_z as the basis of the matrix representation of the Hamiltonian (12). This is possible since the Hamiltonian (12) commutes with I_z :

$$[\mathcal{H}, I_z] = 0, \quad (13)$$

so that both $\mathcal{H}$ and I_z have the common set of eigenvectors. In general the dimensionality of the matrix representation of the Hamiltonian is $2^N \times 2^N$.

As usual, we write the eigenvectors of the operator I_z in terms of the Dirac notations. Let

$$|n_1 \dots n_N\rangle \quad (14)$$

be the eigenvector of the operator I_z where the i th spin is directed opposite to the external magnetic field if $n_i = 1$ and along the field if $n_i = 0$. Then the matrix representation H of the Hamiltonian $\mathcal{H}$ gets the following diagonal block structure:

$$H = \text{diag}(H_0, H_1, H_2, \dots, H_N), \quad (15)$$

where the block H_i is associated with the states having i spins directed opposite to the field. The dimensionality of this block is $C_N^i \times C_N^i$.

It is important that in order to study the problem of the single quantum state transfer along the spin 1/2 chain with Hamiltonian (12) only the blocks H_0 and H_1 are needed,

$$\begin{aligned} H_0 &= -\frac{1}{2}(\tilde{\gamma} - N\omega(t)), \\ H_1 &= \frac{1}{2}[D - (\tilde{\gamma} - (N-2)\omega(t))I], \end{aligned} \quad (16)$$

$$D = \begin{pmatrix} A_{11} & D_1 & D_{13} & \cdots & D_{1(N-2)} & D_{1(N-1)} & D_{1N} \\ D_1 & A_{22} & D_2 & \cdots & D_{2(N-2)} & D_{2(N-1)} & D_{2N} \\ D_{13} & D_2 & A_{33} & \cdots & D_{3(N-3)} & D_{3(N-1)} & D_{3N} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ D_{1(N-2)} & D_{2(N-2)} & D_{3(N-2)} & \cdots & A_{(N-2)(N-2)} & D_{j-1} & D_{(N-2)N} \\ D_{1(N-1)} & D_{2(N-1)} & D_{3(N-1)} & \cdots & D_{j-1} & A_{(N-1)(N-1)} & D_j \\ D_{1N} & D_{2N} & D_{3N} & \cdots & D_{(N-2)N} & D_j & A_{NN} \end{pmatrix},$$

$$A_{nn} = 2 \sum_{\substack{i=1 \\ i \neq n}}^N D_{in}, \quad \tilde{\gamma} = \sum_{\substack{i,j=1 \\ i < j}}^N D_{ij},$$

where I is $N \times N$ identity matrix. Thus, instead of $2^N \times 2^N$ matrix representation of the Hamiltonian we have $N \times N$ block H_1 and scalar block H_0 , which significantly reduces all calculations.

It was shown [2] that effectiveness of the quantum communication channel may be measured by the fidelities of the state transfers between p_n th and p_m th nodes:

$$F_{p_n p_m}^{(N)}(t) = \frac{|f_{p_n p_m}^{(N)}(t)| \cos \Gamma_{p_n p_m}^{(N)}(t)}{3} + \frac{|f_{p_n p_m}^{(N)}(t)|^2}{6} + \frac{1}{2}, \quad (17)$$

where the amplitudes $f_{p_n p_m}^{(N)}$ and the phases $\Gamma_{p_n p_m}^{(N)}$ are defined as follows ($n, m = 1, \dots, \mathcal{N}$):

$$f_{p_n p_m}^{(N)}(t) = \langle p_m | e^{-i\mathcal{H}t} | p_n \rangle = \sum_{j=1}^N u_{p_n j} u_{p_m j} e^{-i\lambda_j t/2} e^{\frac{i}{2} \left(\gamma t - (N-2) \int_0^t \omega(\tau) d\tau \right)}, \quad (18)$$

$$\Gamma_{p_n p_m}^{(N)}(t) = \varphi_{p_n p_m}^{(N)}(t) - \int_0^t \omega(\tau) d\tau, \quad (19)$$

$$\varphi_{p_n p_m}^{(N)}(t) = \arg \left(\sum_{j=1}^N u_{p_n j} u_{p_m j} e^{-i\lambda_j t/2} \right). \quad (20)$$

Here u_{ij} , $i, j = 1, \dots, N$, are components of the eigenvector u_j corresponding to the eigenvalue λ_j of the matrix D : $Du_j = \lambda_j u_j$. The effectiveness of the state transfer is characterized by the set of parameters $\bar{F}_{p_n p_m}^{(N)} \equiv F_{p_n p_m}^{(N)}(t_{p_n p_m}^{(N)})$, $n, m = 1, \dots, \mathcal{N}$. It is evident, that these parameters take maximum values if

$$\bar{\Gamma}_{p_n p_m}^{(N)} \equiv \Gamma_{p_n p_m}^{(N)}(\bar{t}_{p_n p_m}^{(N)}) = 0 \pmod{2\pi}, \quad n, m = 1, \dots, \mathcal{N}. \quad (21)$$

which may be considered as the system of equation defining (not uniquely) the time dependence $\omega(t)$ (which must be positive over the interval $0 \leq t \leq T_{\mathcal{N}}^{(N)}$):

$$\bar{\omega}_{p_n p_m}^{(N)} \equiv \int_0^{\bar{t}_{p_n p_m}^{(N)}} \omega(\tau) d\tau = \bar{\varphi}_{p_n p_m}^{(N)} \equiv \varphi_{p_n p_m}^{(N)}(\bar{t}_{p_n p_m}^{(N)}) \pmod{2\pi}, \quad (22)$$

$n, m = 1, \dots, \mathcal{N}.$

Then the values of the fidelities $F_{p_n p_m}^{(N)}(\bar{t}_{p_n p_m}^{(N)})$ are defined by $|f_{p_n p_m}^{(N)}(\bar{t}_{p_n p_m}^{(N)})|$ or, equivalently, by the probabilities $\bar{P}_{p_n p_m}^{(N)}$. For this reason, namely probabilities $\bar{P}_{p_n p_m}^{(N)}$ will be studied in the rest of this paper. Phases $\varphi_{p_n p_m}^{(N)}(t)$ (which are independent on $\omega(t)$) will be taken as parameters of the HPST($N; \mathcal{L}$) instead of the parameters $\bar{\varphi}_{p_n p_m}^{(N)}$ in the set (8), see Tables 1 - 6. Remark that, deriving eqs.(17-20) we took into account that the energy of the ground state E_0 is not zero: $E_0 = \hbar H_0$.

An explicite example of the function $\omega(t)$ constructed in accordance with eqs.(22) with $N = \mathcal{N} = 4$ will be given in Sec.3, see Example 1.

3 Simplest chains allowing the HPSTs among the four nodes.

The HPSTs from the end node to (some of) the inner nodes of the spin chain becomes possible due to the following mechanism. Let us take two identical chains $L_1(N_1)$ which allow the HPST between their end nodes. Let $\bar{D}$ be the minimal coupling constant between the nearest neighbours in these chains. We connect these chains by the weak bond with the coupling constant $D_{N_1} \ll \bar{D}$, resulting in the chain $L_{11}(N_1, 0, N_1)$ of $N = 2N_1$ nodes, see Fig.4. Then




Fig. 4. Spin 1/2 chain $L_{11}(N_1, 0, N_1)$ with the HPSTs among the four nodes p_i , $i = 1, 2, 3, 4$, i.e. $\mathcal{L} = \{p_i, i = 1, 2, 3, 4\}$, $\mathcal{N} = 4$; C_1 is absent.

$$\mathcal{L} = \{\text{end nodes of chains } L_1(N_1)\} = \{p_1, p_2, p_3, p_4\}, \quad \mathcal{N} = 4. \quad (23)$$

These HPSTs may be understood as follows. Let spin p_1 be directed opposite to the external magnetic field while all other spins of the chain $L_{11}(N_1, 0, N_1)$ be directed along the field initially. Due to the small coupling constant D_{N_1} this excited state remains inside of the first chain $L_1(N_1)$ for the long time with the high probability to be detected in either p_1 or p_2 (since, by definition, the chain $L_1(N_1)$ provides HPST between end nodes). However, due to the bond between two chains $L_1(N_1)$, the excited state will be transferred to the second chain $L_1(N_1)$ after comparatively long time interval T_{tr} with the high probability. If this happens, then the excited state remains in this chain for the long time with high probability to be detected in either p_3 or p_4 . Similarly, after one more time interval T_{tr} the excited state will return to the first chain $L_1(N_1)$ with the high probability, and so on. Parameter D_{N_1} may be fixed by the two conditions (9) and (10). Hereafter in this section we take $P_0 = 0.9$ in the definition (7). Consider two simple examples.

Example 1: the HPSTs in the chain $L_{11}(2, 0, 2)$. The simplest example of the chain $L_1(N_1)$ providing the ideal end-to-end state transfer is the chain of two nodes, i.e. $N_1 = 2$. Connecting two equivalent chains $L_1(N_1)$ we obtain the chain $L_{11}(2, 0, 2)$ with $N = \mathcal{N} = 4$, see Fig.5.




Fig. 5. Spin 1/2 chain $L_{11}(2,0,2)$ with the HPSTs among all nodes $p_i \equiv i$, $i = 1, 2, 3, 4$, i.e. $\mathcal{L} = \{1, 2, 3, 4\}$, $N=4$. This figure is equivalent to Fig.4 with $N_1 = 2$. The optimal value of the parameter δ is 0.196.

In this case the set $\mathcal{L}$ consists of all nodes of the chain $L_{11}(2,0,2)$. We set $D_1 = D_3 = 1$ for simplicity and vary $D_2 = \delta$ in order to satisfy the conditions (9) and (10). We have found that the optimal parameters of the HPST(4;1,2,3,4) correspond to $\delta = 0.196$. These parameters are represented in Table 1, see also Fig.11.

Let us find the time evolution of the external magnetic field which satisfies conditions (22), i.e such $\omega(t)$ that

$$\bar{\omega}_{nm}^{(4)} \equiv \int_0^{\bar{t}_{nm}^{(4)}} \omega(\tau) d\tau = \bar{\varphi}_{nm}^{(4)} \mod(2\pi), \quad n, m = 1, 2, 3, 4. \quad (24)$$

We may write conditions (24) as the following system of five equations:

$$\begin{aligned} \bar{\omega}_{12}^{(4)} &= 0, \quad \bar{\omega}_{12}^{(4)} = \bar{\varphi}_{12}^{(4)}, \quad \bar{\omega}_{13}^{(4)} = \bar{\varphi}_{13}^{(4)} + 8\pi, \\ \bar{\omega}_{14}^{(4)} &= \bar{\varphi}_{14}^{(4)} + 8\pi, \quad \bar{\omega}_{23}^{(4)} = \bar{\varphi}_{23}^{(4)} + 6\pi, \end{aligned} \quad (25)$$

which is satisfied by the following function $\omega(t)$:

$$\begin{aligned} \int_0^t \omega(\tau) d\tau &= a_4 t^4 + a_3 t^3 + a_2 t^2 + a_1 t \Rightarrow \\ \omega(t) &= 4a_4 t^3 + 3a_3 t^2 + 2a_2 t + a_1, \\ a_1 &= 4.1241 \times 10^{-1}, \quad a_2 = -6.9297 \times 10^{-3}, \\ a_3 &= 2.3114 \times 10^{-4}, \quad a_4 = -1.9971 \times 10^{-6}. \end{aligned} \quad (26)$$

We see that the function $\omega(t)$ constructed in this way is positive over the interval $0 \leq t \leq \bar{t}_{14}^{(4)} = \max_{n,m=1,\dots,4} (\bar{t}_{nm}^{(4)})$.

Example 2: the HPSTs in the chain $L_{11}(3, 0, 3)$. We take two homogeneous chains of three nodes $L_1(3)$. It is known that the ideal state transfer is possible between the end nodes of these chains [3]. We connect them by the weak bond obtaining the chain shown in Fig.6. Thus, $\mathcal{L} = 1, 3, 4, 6$. Let




Fig. 6. Spin 1/2 chain $L_{11}(3, 0, 3)$ with the HPSTs among the four nodes $p_1 = 1$, $p_2 = 3$, $p_3 = 4$, $p_4 = 6$; $\mathcal{L} = \{1, 3, 4, 6\}$, $\mathcal{N} = 4$. $\delta_1 = 1$, $\delta_2 = 0.028$ in Example 2 of Sec.3 and $\delta_1 = 0.769$, $\delta_2 = 0.092$ in the chain $\hat{L}_{11}(3, 0, 3)$ considered in Example 1 of Sec.5

$D_1 = D_2 = D_4 = D_5 = 1$, $D_3 = \delta_2$ in this chain. We vary δ_2 to obtain the best correspondence to the conditions (9) and (10). The optimal value is $\delta_2 = 0.028$. The appropriate parameters of the HPST(6;1,3,4,6) are represented in Table 2.

3.1 Modification of the chain $L_{11}(N_1, 0, N_1)$

The chain shown in Fig.4 is convenient for the state transfer between two chains L_1 if only the distance between them (i.e. between p_2 and p_3) is not too long. Otherwise the time interval $T_4^{(2N_1)}$ becomes very long. To decrease this time interval we suggest the following modification of the chain $L_{11}(N_1, 0, N_1)$.

Let us take a symmetrical chain $C_1(M_1)$ of M_1 nodes with maximal coupling constant between neighbours $\mathcal{D}$ satisfying the following condition: $\mathcal{D} < \bar{D}$. Using the coupling constant $\mathcal{D}_{12} \ll \bar{D}$ we may construct the chain $L_{11}(N_1, M_1, N_1)$ of $N = 2N_1 + M_1$ nodes shown in Fig.7. Here $\mathcal{L}$ consists of the end nodes of chains L_1 and may not involve any node of $C_1(M_1)$. This statement is valid due to the fact, that the probability for the spin to be detected in the chain $C_1(M_1)$ may not be high because of the small coupling constants both inside of this chain and $\mathcal{D}_{12}$. These coupling constants should be fixed by the conditions (9) and (10).

Example: the HPSTs in the chain $L_{11}(2, 2, 2)$. We consider the HPST(6;1,2,5,6) in the chain $L_{11}(2, 2, 2)$ shown in Fig.8.

This is a chain of 6 nodes ($N = 6$). We fix $D_1 = D_5 = 1$ and vary parameters




Fig. 7. Spin 1/2 chain $L_{11}(N_1, M_1, N_1)$ with the HPSTs among the four nodes p_i , $i = 1, 2, 3, 4$; $\mathcal{L} = \{p_1, p_2, p_3, p_4\}$, $\mathcal{N} = 4$. Chain $C_1(M_1)$ is introduced to decrease the parameter $T_4^{(N)}$, $N = 2N_1 + M_1$ (compare with the scheme in Fig.4).




Fig. 8. Spin 1/2 chain $L_{11}(2, 2, 2)$ with the HPSTs among the four nodes $p_1 = 1$, $p_2 = 2$, $p_3 = 5$, $p_4 = 6$; $\mathcal{L} = \{1, 2, 5, 6\}$, $\mathcal{N} = 4$. Chain $C_1(2)$ is introduced to decrease the parameter $T_4^{(6)}$ (compare with the scheme in Fig.7). Optimal parameters of the HPST(6;1,2,5,6) correspond to $\delta_1 = 0.224$, $\delta_2 = 0.649$.

$D_2 = D_4 = \delta_1$ and $D_3 = \delta_2$ with the purpose to decrease the parameter $T_4^{(6)}$ and satisfy the conditions (9,10). We have found that the following values of the parameters δ_i yield a good result: $\delta_1 = 0.224$ and $\delta_2 = 0.649$. The appropriate parameters of the HPST(6;1,2,5,6) are represented in Table 3.

Here we demonstrate that the intermediate chain $C_1(2)$ in the chain $L_{11}(2, 2, 2)$ speeds up the state transfer between p_2 and p_3 separated by the distance $R = 2(\Gamma/\delta_1)^{1/3} + (\Gamma/\delta_2)^{1/3}$. For this purpose we compare the parameters $\bar{t}_{23}^{(4)}$ for the chain $L_{11}(2, 0, 2)$ (see Fig.5) with $D_2 = \Gamma/R^3 \approx 0.011$ and parameter $\bar{t}_{25}^{(6)} \approx 64.4$ from the Table 3. Numerical simulation shows that $\bar{t}_{23}^{(4)} \approx 2000$, i.e. $\bar{t}_{23}^{(4)}/\bar{t}_{25}^{(6)} \approx 31$.

4 Simplest chains with the HPSTs among eight nodes.

Chains $L_{11}(N_1, 0, N_1)$ and $L_{11}(N_1, M_1, N_1)$ considered in Sec.3 provide the HPSTs between any two nodes out of the set $\mathcal{L}$ consisting of four nodes. However, the number of such nodes may be increased using the following obvious generalization of these chains.

Let us take two chains $L_{12}(N_1, M_1, N_1)$ and $L_{21}(N_1, M_1, N_1)$ (see Fig.1) and the symmetrical chain $C_2(M_2)$. The maximal coupling constant $\bar{D}$ between the nearest neighbours in $C_2(M_2)$ must satisfy inequality $\bar{D} < \min(\mathcal{D}_{12}, \mathcal{D}_{13}, \bar{D})$, where $\bar{D}$ is the minimal coupling constant between the nearest neighbours in $L_i(N_1)$, $i = 1, 2$. Using the coupling constant $\mathcal{D}_{24}$, $\mathcal{D}_{24} \ll \bar{D}$, we construct the chain of $N = 4N_1 + 2M_1 + M_2$ nodes shown in Fig.9. Here $\mathcal{L}$ consists of the




Fig. 9. Spin 1/2 chain $L_{1221}(N_1, M_1, N_1, M_2, N_1, M_1, N_1)$ with the HPSTs among the eight nodes p_i , $i = 1, \dots, 8$; $\mathcal{L} = \{p_i, i = 1, \dots, 8\}$, $\mathcal{N} = 8$.

eight end nodes of chains L_1 and L_2 : $\mathcal{L} = \{p_i, i = 1, \dots, 8\}$, $\mathcal{N} = 8$.

It is obvious that this algorithm may be extended to construct chains with $\mathcal{L}$ consisting of 2^s nodes $s = 1, 2, \dots$. Consider the simplest example where we take $P_0 = 0.8$ (see eq.(7)).

Example: the HPSTs in the chain $L_{1111}(2, 0, 2, 0, 2, 0, 2)$. We consider the chain of eight nodes as one obtained by joining of two 4-nodes chains $L_{11}(2, 0, 2)$ constructed in Example 1 of Sec.3, see Fig.10 where $\delta_2 = 1$.




Fig. 10. Spin 1/2 chain $L_{1221}(2, 0, 2, 0, 2, 0, 2)$ with the HPSTs among the eight nodes $p_i = i$, $i = 1, \dots, 8$; $\mathcal{L} = \{1, \dots, 8\}$, $\mathcal{N} = 8$; $\delta_1 = 0.196$, $\delta_2 = 1$, $\delta_3 = 0.010$ in the chain $L_{1111}(2, 0, 2, 0, 2, 0, 2)$ considered in Example of Sec.4 ($L_2(2) \equiv L_1(2)$ in this case) and $\delta_1 = 0.247$, $\delta_2 = 0.977$, $\delta_3 = 0.018$ in the chain $\hat{L}_{1221}(2, 0, 2, 0, 2, 0, 2)$ considered in Example 2 of Sec.5.

To provide the HPSTs among all nodes we must take a small coupling constant D_4 between these two chains. Namely, it must be less than $\delta = 0.196$ introduced in Example 1 of Sec.3. We take $D_1 = D_7 = 1$, $D_2 = D_6 = \delta_1 = 0.196$, $D_3 = D_5 = \delta_2 = 1$, $D_4 = \delta_3 \ll 0.196$. The disadvantages of this chain are big parameter $T_8^{(8)}$ and comparatively small values of $P_{ij}^{(8)}$, $i, j = 1, \dots, 8$ ($i \neq j$). After optimization we obtain $\delta_3 = 0.010$. The parameters of the HPST(8;1,2,3,4,5,6,7,8) are represented in Table 4.

5 Deformations of the chains improving the parameter $T_{\mathcal{N}}^{(N)}$

It may be shown that the parameters of the HPSTs may be improved varying all coupling constants in the chains $L_{11}(N_1, M_1, N_1)$ and $L_{1221}(N_1, M_1, N_1, M_2, N_1, M_1, N_1)$ in a proper way, i.e. varying not only the coupling constants between chains $L_1(N_1)$ (like it was done in Secs.3 and 4) but also the coupling constants inside of $L_1(N_1)$. This allows one to decrease significantly parameter $T_{\mathcal{N}}^{(N)}$. After such process the above chains will be reduced to the "optimized" chains which we call $\hat{L}_{11}(N_1, M_1, N_1)$ and $\hat{L}_{1221}(N_1, M_1, N_1, M_2, N_1, M_1, N_1)$ respectively (compare the parameters from Tables 2 and 5 and from Tables 4 and 6).

Example 1: the HPSTs in the chain $\hat{L}_{11}(3, 0, 3)$, $P_0 = 0.9$. The purpose of this section is to decrease the parameter $T_4^{(6)}$ which have been found in Example 2 of Sec.3, see Table 2 and Fig.6. We take $D_1 = D_5 = 1$, $D_2 = D_4 = \delta_1$, $D_3 = \delta_2$ and vary δ_i , $i = 1, 2$ keeping in mind the conditions (9,10). We have found that the optimal parameters of the HPST(6;1,3,4,6) correspond to the $\delta_1 = 0.769$ and $\delta_2 = 0.092$, see Table 5 and Fig.6.

Example 2: the HPSTs in the chain $\hat{L}_{1221}(2, 0, 2, 0, 2, 0, 2)$, $P_0 = 0.8$. The parameter $T_8^{(8)}$ obtained in Example 1 of Sec.4 may be decreased varying the coupling constants in the chain, see Fig.10. Thus, the optimal parameters of the HPST(8;1,2,3,4,5,6,7,8) have been found for $\delta_1 = 0.247$, $\delta_2 = 0.977$, $\delta_3 = 0.018$, see Table 6 and Fig.10.

$p_i \backslash p_j$	1	2	3	4
1		0.931 3.040 1.196	0.904 55.533 -3.010	0.912 58.585 -1.806
2	0.931 3.040 1.196		0.906 52.548 1.998	0.904 55.533 -3.010
3	0.904 55.533 -3.010	0.906 52.548 1.998		0.931 3.040 1.196
4	0.912 58.585 -1.806	0.904 55.533 -3.010	0.931 3.040 1.196	

Table 1

The parameters $\bar{P}_{ij}^{(4)}$ (the first number in the box), $\bar{t}_{ij}^{(4)}$ (the second number in the box) and $\bar{\varphi}_{ij}^{(4)}$ (the third number in the box) of the HPST(4;1,2,3,4) in $L_{11}(2, 0, 2)$




Fig. 11. The time dependence of the probabilities $P_{1i}^{(4)}$, $i = 1, 2, 3, 4$ and $P_{23}^{(4)}$ in the chain $L_{11}(2, 0, 2)$. The marked points correspond to the parameters $\bar{P}_{ij}^{(4)}$ and $\bar{t}_{ij}^{(4)}$ of the HPST(4;1,2,3,4), see also Table 1.

$p_i \backslash p_j$	1	3	4	6
1		0.978 11.595 3.120	0.909 426.354 0.313	0.927 414.760 -2.812
3	0.978 11.595 3.120		0.919 414.762 -2.812	0.909 426.354 0.313
4	0.909 426.354 0.313	0.919 414.762 -2.812		0.978 11.595 3.120
6	0.927 414.760 -2.812	0.909 426.354 0.313	0.978 11.595 3.120	

Table 2

The parameters $\bar{P}_{p_i p_j}^{(6)}$ (the first number in the box), $\bar{t}_{p_i p_j}^{(6)}$ (the second number in the box) and $\bar{\varphi}_{p_i p_j}^{(6)}$ (the third number in the box) of the HPST(6;1,3,4,6) in $L_{11}(3, 0, 3)$

$p_i \backslash p_j$	1	2	5	6
1		0.913 3.005 1.147	0.926 67.364 2.120	0.971 70.375 -3.003
2	0.913 3.005 1.147		0.934 64.400 0.925	0.926 67.364 2.120
5	0.926 67.364 2.120	0.934 64.400 0.925		0.913 3.005 1.147
6	0.971 70.375 -3.003	0.926 67.364 2.120	0.913 3.005 1.147	

Table 3

The parameters $\bar{P}_{p_i p_j}^{(6)}$ ((the first number in the box), $\bar{t}_{p_i p_j}^{(6)}$ (the second number in the box) and $\bar{\varphi}_{p_i p_j}^{(6)}$ (the third number in the box) of the HPST(6;1,2,5,6) in $L_{12}^C(2, 2, 2)$

6 Conclusions

Using the numerical simulations we demonstrate that the N -nodes spin 1/2 chains with wide spread of the properly adjusted coupling constants allow the HPSTs between different nodes. It is important that the whole Hamiltonian (12) rather than approximation by the nearest neighbour interaction must be used for the correct description of the HPSTs in such chains.

We have found that two spin 1/2 chains $L_1(N_1)$ with the HPST between end nodes may be connected by a relatively weak bond to get a chain with the HPSTs among four nodes, see Sec.3. In turn, having two such chains we may connect them by another weak bond to get a chain with the HPSTs among eight nodes (see Sec.4), and so on. Formally, the number of the nodes allowing the HPSTs among all of them may be 2^s , where $s = 1, 2, \dots$. However, the disadvantage of such chains is a rapid increase of the time interval $T_{\mathcal{N}}^{(N)}$ with the number of nodes involved in the HPSTs. The mechanism decreasing $T_{\mathcal{N}}^{(N)}$ would be important for the implementation of these chains.

We also demonstrate that the speedup of the state transfer between the nodes p_i and p_{i+1} separated by the distance R may be achieved using the intermediate chain with properly adjusted coupling constants, see Sec.3.1.

This work is supported by Russian Foundation for Basic Research through the grant 07-07-00048 and by the Program of the Department of Chemistry and Material Science of RAS No.18.

$p_i \backslash p_j$	1	2	3	4	5	6	7	8
1		0.930 3.040 1.181	0.882 55.557 2.666	0.854 52.463 1.573	0.813 3385.361 -2.762	0.835 3382.303 2.367	0.818 3329.745 0.947	0.862 3326.706 -0.245
2	0.930 3.040 1.181		0.871 52.563 1.431	0.855 55.556 2.666	0.812 3382.302 2.367	0.829 3379.311 1.141	0.886 3326.702 -0.227	0.818 3329.745 0.947
3	0.882 55.557 2.666	0.871 52.563 1.431		0.933 3.044 1.143	0.830 3329.851 0.754	0.893 3326.800 -0.376	0.829 3379.311 1.141	0.835 3382.303 2.367
4	0.854 52.463 1.573	0.855 55.556 2.666	0.933 3.044 1.143		0.876 3326.807 -0.396	0.830 3329.851 0.754	0.812 3382.302 2.367	0.813 3385.361 -2.762
5	0.813 3385.361 -2.762	0.812 3382.302 2.367	0.830 3329.851 0.754	0.876 3326.807 -0.396		0.933 3.044 1.143	0.855 55.556 2.666	0.854 52.463 1.573
6	0.835 3382.303 2.367	0.829 3379.311 1.141	0.893 3326.800 -0.376	0.830 3329.851 0.754	0.933 3.044 1.143		0.871 52.563 1.431	0.882 55.557 2.666
7	0.818 3329.745 0.947	0.886 3326.702 -0.227	0.829 3379.311 1.141	0.812 3382.302 2.367	0.855 55.556 2.666	0.871 52.563 1.431		0.930 3.040 1.181
8	0.862 3326.706 -0.245	0.818 3329.745 0.947	0.835 3382.303 2.367	0.813 3385.361 -2.762	0.854 52.463 1.573	0.882 55.557 2.666	0.930 3.040 1.181	

Table 4

The parameters $\bar{P}_{p_i p_j}^{(8)}$ (the first number in the box), $\bar{t}_{p_i p_j}^{(8)}$ (the second number in the box) and $\bar{\varphi}_{p_i p_j}^{(8)}$ (the third number in the box) of the HPST(8;1,2,3,4,5,6,7,8) in $L_{1111}^C(2, 0, 2, 0, 2, 0, 2)$

References

- [1] E.I.Kuznetsova and A.I.Zenchuk, Phys.Lett.A, **372** (2008) 6134
- [2] S.Bose, Phys.Rev.Lett, **91** (2003) 207901
- [3] M.Christandl, N.Datta, A.Ekert and A.J.Landahl, Phys.Rev.Lett. **92** (2004) 187902
- [4] E.B.Fel'dman, R.Brüschweiler and R.R.Ernst, Chem.Phys.Lett. **294** (1998) 297
- [5] C.Albanese, M.Christandle, N.Datta and A.Ekert, Phys.Rev.Lett., **93**, 230502 (2004)

$p_i \backslash p_j$	1	3	4	6
1		0.908 13.095 2.588	0.919 87.366 1.483	0.916 74.231 -1.044
3	0.908 13.095 2.588		0.905 74.324 -1.178	0.919 87.366 1.483
4	0.919 87.366 1.483	0.905 74.324 -1.178		0.908 13.095 2.588
6	0.916 74.231 -1.044	0.919 87.366 1.483	0.908 13.095 2.588	

Table 5

The parameters $\bar{P}_{p_i p_j}^{(6)}$ (the first number in the box), $\bar{t}_{p_i p_j}^{(6)}$ (the second number in the box) and $\bar{\varphi}_{p_i p_j}^{(6)}$ (the third number in the box) of the HPST(6;1,3,4,6) in $\hat{L}_{11}(3, 0, 3)$

- [6] P.Karbach and J.Stolze, Phys.Rev.A, **72**, 030301(R) (2005)
- [7] E.B.Fel'dman and M.G.Rudavets, JETP Letters, **81** (2005) 47
- [8] E.I.Kuznetsova and E.B.Fel'dman, J.Exp.Theor.Phys., **102** (2006) 882
- [9] L.C.Venuti, S.M.Giampaolo, F.Illuminati and P.Zanardi, Phys.Rev.A **76** (2007) 052328
- [10] G.Gualdi, V.Kostak, I.Marzoli and P.Tombesi, arXiv:0807.5018 [quant-ph]
- [11] G.B.Furman, V.M.Meerovich and V.L.Sokolovsky, Phys.Rev.A, **77**, 062330 (2008)
- [12] Abragam A. The principles of nuclear magnetism, Oxford, Clarendon Press, 1961

$p_i \backslash p_j$	1	2	3	4	5	6	7	8
1		0.893 2.984 1.139	0.858 43.271 -3.070	0.883 46.305 -1.954	0.847 1537.838 -1.667	0.862 1534.792 -2.751	0.812 1488.470 -0.718	0.870 1491.478 0.383
2	0.893 2.984 1.139		0.873 46.284 -1.976	0.820 43.267 -3.067	0.823 1534.790 -2.749	0.866 1537.766 -1.600	0.906 1491.433 0.464	0.812 1488.470 -0.718
3	0.858 43.271 -3.070	0.873 46.284 -1.976		0.895 3.059 1.064	0.814 1488.517 -0.836	0.901 1491.551 0.276	0.866 1537.766 -1.600	0.862 1534.792 -2.751
4	0.883 46.305 -1.954	0.820 43.267 -3.067	0.895 3.059 1.064		0.871 1491.604 0.187	0.814 1488.517 -0.836	0.823 1534.790 -2.749	0.847 1537.838 -1.667
5	0.847 1537.838 -1.667	0.823 1534.790 -2.749	0.814 1488.517 -0.836	0.871 1491.604 0.187		0.895 3.059 1.064	0.820 43.267 -3.067	0.883 46.305 -1.954
6	0.862 1534.792 -2.751	0.866 1537.766 -1.600	0.901 1491.551 0.276	0.814 1488.517 -0.836	0.895 3.059 1.064		0.873 46.284 -1.976	0.858 43.271 -3.070
7	0.812 1488.470 -0.718	0.906 1491.433 0.464	0.866 1537.766 -1.600	0.823 1534.790 -2.749	0.820 43.267 -3.067	0.873 46.284 -1.976		0.893 2.984 1.139
8	0.870 1491.478 0.383	0.812 1488.470 -0.718	0.862 1534.792 -2.751	0.847 1537.838 -1.667	0.883 46.305 -1.954	0.858 43.271 -3.070	0.893 2.984 1.139	

Table 6

The parameters $\bar{P}_{p_i p_j}^{(8)}$ (the first number in the box), $\bar{t}_{p_i p_j}^{(8)}$ (the second number in the box) and $\bar{\varphi}_{p_i p_j}^{(8)}$ (the third number in the box) of the HPST(8;1,2,3,4,5,6,7,8) in $\hat{L}_{1221}^C(2, 0, 2, 0, 2, 0, 2)$